

Relating bars with the environment in the nearby Universe

Héctor J. Martínez^{*} & Hernán Muriel

Instituto de Astronomía Teórica y Experimental, IATE, CONICET–Observatorio Astronómico, Universidad Nacional de Córdoba, Laprida 854, X5000BGR, Córdoba, Argentina

29 September 2011

ABSTRACT

We study the correlation between the fraction of barred spiral galaxies and environmental parameters of galaxies to understand in which environments the bars are more commonly found. For this purpose we apply the Blanton et al. technique to a sample of spiral galaxies drawn from the Nair & Abraham catalogue. Our results agree with previous findings in which the fraction of barred galaxies is almost insensitive to environment.

Key words: galaxies: fundamental parameters – galaxies: evolution – galaxies: statistics

1 INTRODUCTION

Bars are believed to be closely related with the dynamical evolution of disk galaxies and to play an important role in redistributing the angular momentum between dark and baryonic matter (Weinberg 1985; Debattista & Sellwood 1998). Athanassoula (2003) proposed that this exchange of angular momentum is closely related to the density and velocity dispersion of the host halo. Other roles that have been assigned to bars are: to transport material to the centre and ignite starbursts (Sheth et al. 2005) and/or feed the central black hole (Shlosman et al. 1990; Corsini et al. 2003), however no direct evidence of this is seen (Mulchaey & Regan 1997); change chemical abundance gradient (Zaritsky 1992; Martin & Roy 1994); bars can trigger star formation along themselves, or have a lot of gas and no star formation (Kenney & Lord 1991; Sheth et al. 2002). Bars are also associated with circumnuclear star formation activity (Sérsic & Pastoriza 1967; Ho et al. 1997; Sheth et al. 2000); the formation of nuclear pseudo-bulges (Kormendy 1982; Kormendy & Kennicutt 2004), rings (Schwarz 1981; Buta 1986; Buta & Combes 1996) and, possibly, spiral arms (Lindblad 1960; Elmegreen & Elmegreen 1985).

For a fixed circular velocity, barred and unbarred galaxies have similar properties like luminosity, scale lengths, star formation and colour (Courteau et al. 2003), which does not necessary imply they have followed the same evolutionary paths (see Sheth et al. 2008). de Vaucouleurs (1963) using blue plates found that more than 60% of nearby galaxies are barred. Similar fractions are observed in the near-infrared (Eskridge et al. 2000; Laurikainen et al. 2004; Menéndez-Delmestre et al. 2007). In numerical simulations, bars appear naturally once the cold and rotationally supported disk is in place (see Athanassoula 2005 for a review), although, the precise mechanisms that drive this phenomenon are not well established yet. Among the internal mechanisms that can produce

bars is the instability of the disk (Sellwood & Wilkinson 1993; Heller et al. 2007; Athanassoula 2008).

It has also been suggested that environment could also be important in leading to the formation of bars, although results are contradictory. In some numerical simulations bars are created in mergers and interactions (Walker et al. 1996; Mihos et al. 1997; Berentzen et al. 2004), it is transient in others (Gerin et al. 1990) and in some others bars are destroyed as the galaxy becomes an elliptical. Elmegreen et al. (1990) analyse galaxies in binary and group systems and in the field and find a correlation between the bar fraction and environment for early-type spirals, with the highest fraction corresponding to binary systems. On the other hand, van den Bergh (2002) use 930 galaxies from the Palomar Sky Survey and conclude that the bar fraction does not depend on the environment. An environment that has been particularly studied is that of galaxy clusters. Méndez-Abreu et al. (2010) study both, the centre and the infall regions of the Coma Cluster, in a wide range of magnitudes. They find that bars are hosted in galaxies in a tight interval of masses and luminosities ($10^9 \leq M_*/M_\odot \leq 10^{11}$ and $-22 < M_r < -17$ respectively). These authors do not find a significant difference in the fraction of bars between galaxies in the centre and in the infall regions suggesting that the cluster environment plays a second-order role in the bar formation/evolution.

The purpose of this paper is to evaluate different parameters that characterise the environment as possible generators of bars. We use the technique proposed by Blanton et al. (2005) and significance criteria introduced by Martínez et al. (2008) to samples of galaxies drawn from the catalogue by Nair & Abraham (2010a) (hereafter NA10). The paper is organised as follows: section 2 describe the sample of galaxies and the environmental properties we use throughout our work; section 3 presents the results; finally we provide a discussion of our findings in section 4. Throughout this paper we assume a flat cosmological model with parameters $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$.

^{*} E-mail: julian@oac.uncor.edu

2 THE GALAXY SAMPLE

For the purposes of this paper we use a sample of galaxies drawn from the catalogue by NA10. This catalogue presents detailed visual classification for 14034 galaxies in the Main Galaxy Sample (MGS; Strauss et al. 2002) of the Fourth Data Release of the SDSS (DR4; Adelman-McCarthy et al. 2006) that constitute a complete sample with $0.01 \leq z \leq 0.1$ and down to a limiting extinction corrected apparent magnitude of $g = 16$. Each galaxy in the catalogue has been morphologically classified by NA10 into T-Types. Additionally, they recorded the existence of structures such as bars, rings, lenses, tails, warps, etc.

In our analyses below, we include all galaxies classified as spirals (T-Type ≥ 0) in the NA10 catalogue which have axial ratio $b/a > 0.55$. Below this cut-off the fraction of barred galaxies drops dramatically (fig. 21 in NA10). There are no other important incompleteness regarding bar identifications in the NA10. Our sample has 5508 galaxies, among which 1841 are barred.

For the analysis of the correlation between the existence of bars and the environment in which galaxies are located, we selected 4 measures of environment quoted in the NA10 catalogue and also computed the projected distance to the nearest MGS neighbour brighter than $M_r = -20$ (that is, a volume limited sample of galaxies up to $z = 0.1$) and with $c|\Delta z| \leq 1000 \text{ km s}^{-1}$ as another environment measure:

- (i) $L_{\text{group},g}$: group luminosity ($9.5 \leq \log(L_{\text{group},g}/L_{\odot,g}) \leq 12.5$) from Yang et al. (2007).
- (ii) M_{group} : group mass ($9.5 \leq \log(M_{\text{group}}/M_{\odot}) \leq 13.0$) from Yang et al. (2007).
- (iii) M_{halo} : group halo mass ($11.5 \leq \log(M_{\text{halo}}/M_{\odot}) \leq 15.5$) from Yang et al. (2007).
- (iv) Σ_5 : 5th neighbour projected density ($-0.8 \leq \log(\Sigma_5/\text{Mpc}^{-2}) \leq 1.8$) from Baldry et al. (2006).
- (v) r_{NN} : projected distance to the nearest neighbour ($0 \leq r_{NN}/\text{kpc} \leq 1500$) computed in this work. We have not introduced corrections accounting for the well known fiber collision incompleteness of the MGS, nevertheless, this should not bias our results since, both, barred and non-barred spirals should be affected in the same way.

All cut-offs in the lists above were imposed as a compromise between probing the largest possible volume in the space of parameters and, at the same time, having enough galaxies per bin to properly carry out the correlation analyses detailed in the next section.

3 RELATING BARS WITH ENVIRONMENTS

We explore the ability of different environment parameters to predict the fraction of barred galaxies by using the σ_X statistics as defined by Blanton et al. (2005). Briefly, given a set of environmental properties, $X_1, \dots, X_N$, the fraction of barred galaxies, f_{bar} , will be, in principle, a function of them: $f_{\text{bar}} = f_{\text{bar}}(X_1, \dots, X_N)$. If we consider now a particular parameter X_I and marginalise f_{bar} over the remaining ones, we get the fraction of barred galaxies as a function of X_I alone, $f_{\text{bar}}(X_I)$. The parameter X_J that correlates best with the fraction of barred galaxies will be the one that minimises the variance σ_{X_J} of $f_{\text{bar}}(X_J)$ after subtracting its global trend (see for details Blanton et al. 2005; Martínez & Muriel 2006; Martínez et al. 2008).

The σ_X statistics provides the environmental parameter that predicts best the presence of bars. This does not mean, however,




Figure 1. The fraction of barred galaxies as a function of different measures of the environment, quoted error-bars are obtained by the bootstrap resampling technique. From upper to lower, panels are sorted according to increasing σ_X values. Shaded areas are the mean values and $\pm 1\sigma$ error-bars from the resamplings. We quote in the bottom the number of galaxies in each bin. Below each panel we show the distributions corresponding to barred (shaded histogram) and non-barred galaxies (empty histogram), both normalised to have the same area.

that the best ranked parameter is a good predictor of bars. To complement the σ_X statistics, we use the significance criterion by Martínez et al. (2008) computed by using bootstrap resamplings.

Since we are dealing with a flux-limited galaxy sample in our analyses, we have weighted each galaxy in our statistics by $1/V_{\max}$ (Schmidt 1968). Given that the fraction of barred galaxies depends on absolute magnitude (Sheth et al. 2008; Nair & Abraham 2010b), and that brighter galaxies tend to be located in higher density environments, the non inclusion of such a weighting scheme can lead to systematics in our analyses.

In Fig. 1 we show the results of our analysis on the relationship between bars and environment. Panels are sorted from top to bottom according to increasing σ_X values. We quote next to each panel the σ_X value along with its significance. Each panel shows the fraction, $f_{\text{bar}}(X)$, of barred galaxies as a function of the corresponding quantity X . Error bars are 1σ bootstrap resampling error-bars. Shaded areas represent the overall mean value of $f_{\text{bar}}(X)$ plus/minus 1σ from the bootstrap resamplings. We also show in the inferior part of each panel the $1/V_{\max}$ weighted histograms of barred (shaded) and non-barred (empty) galaxies, all these histograms are normalised to have the same area. The best predicting parameter is the distance to the nearest neighbour, in the sense that galaxies with bars tend to have closer neighbours. This is followed by the projected galaxy density, the mass of the halo, the group luminosity and finally the group mass. Nevertheless, all the parameters have significance below 68%, i.e., less than 1σ for Gaussian statistics, thus none of them is significant according to our criterion.

Despite that none of the explored parameters are closely correlated to the presence of bars, the trends in Fig. 1 suggest that barred galaxies are slightly more common at higher densities, and in more massive halos. Inspired in the results by Elmegreen et al. (1990), we also check whether the correlation of bars and the environment is stronger for early spirals, by repeating the σ_X computations only for galaxies classified as Sa, Sab and Sb by NA10. We find no evidence of significant correlation for those galaxies either. The results suggest, at best, a second order environmental effect.

We have repeated our computations without the $1/V_{\max}$ weighting scheme, resulting in a different ranking of parameters. In this case we observe that: (i) the overall f_{bar} is slightly higher since f_{bar} is an increasing function of luminosity; (ii) the trends as in Fig. 1 do not qualitatively change, however, (iii) the variances around the global trends, σ_X , do change leading to a different ranking: the group mass ranks first and the remaining four parameters have almost identical σ_X values. This is no surprise since in this case the statistics is dominated by brighter galaxies, that tend to be located in more massive systems, and among them the fraction of bars is higher. Again, we find that significance levels are well below 68%. The non inclusion of a weighting scheme accounting for the fact that we are dealing with an essentially flux limited galaxy sample can lead to wrong conclusions.

Although non-significant according to our criterion, the distance to the nearest neighbour is the environmental parameter that correlates best with the fraction of bars. The nearest neighbour in our analysis is, in all cases, a relatively bright object, $M_r < -20$, since we searched for it in a volume limited (up to $z = 0.1$) sample of galaxies drawn from the MGS to avoid biases with redshift. If the proximity to another galaxy is an important factor for the formation and/or stability of a bar, the mass ratio between a galaxy and its closest neighbour might be important as well. We have further explored the correlation between bars and neighbours in terms of the difference in absolute magnitude between the barred galaxies and their nearest neighbours, which is broadly related to their

mass ratio. We classified spiral galaxies in the NA10 catalogue according to the r -band absolute magnitude difference with the nearest neighbour, $\Delta M = M_{NN} - M$, into two groups: galaxies whose nearest neighbour is brighter or have similar luminosity ($\Delta M \leq 0.5$); and galaxies whose nearest neighbour is fainter ($1.5 > \Delta M > 0.5$). In this case, we searched for the nearest neighbours in the flux limited ($r \leq 17.77$) MGS, and restricted the sample of spiral galaxies in the NA10 to those that are at least 1.5 magnitudes brighter than the absolute magnitude corresponding to the apparent magnitude limit at the galaxy's redshift. For these two subsamples we have studied the trend of the fraction of barred galaxies as a function of the projected distance to the nearest neighbour, as in the *upper panel* of Fig. 1, and have found no significant differences. Thus, according to our results, bars are located in galaxies with no particular preference regarding the luminosity of the nearest neighbour.

4 CONCLUSIONS

We have studied the relationship between the fraction of barred spiral galaxies and a number of environmental parameters in the nearby universe using a complete sample of spiral galaxies taken from the NA10 catalogue. For this purpose we have applied the technique by Blanton et al. (2005) and the significance criterion by Martínez et al. (2008), to a sample of spiral galaxies taken from the NA10 catalogue. Once the range of a set of parameters is defined, this technique measures the ability of each parameter to predict an observable, in this case, the bar fraction.

We have considered in our analysis 5 parameters characterising the environment: group luminosity, group mass, group halo mass (the three of them computed by Yang et al. 2007), the projected galaxy density (computed by Baldry et al. 2006) and the projected distance to the nearest neighbour (computed in this work). Our results indicate that the latter parameter is the one that best predicts the existence of bars, however the signal is only marginal. The small effect is in the sense that spiral galaxies with a nearest neighbour within 0.5 Mpc tend to have a slightly higher fraction of bars. We find no evidence of bars preferring systematically brighter or fainter neighbours. Our finding that the proximity to another galaxy could play a role in the formation of a bar, can be related to the predictions of some numerical simulations in which interactions can trigger bars and/or influence bar properties (e.g. Noguchi 1987; Gerin et al. 1990; Mihos et al. 1997; Miwa & Noguchi 1998; Berentzen et al. 2004). However, due to the low significance we find, our results can also be in agreement with van den Bergh (2002) and Méndez-Abreu et al. (2010), in which the fraction of barred galaxies does not depend on the environment. Numerical simulations have shown that bars form naturally in discs (e.g. Athanassoula 2005 and references therein), thus, bars could be understood in terms of nature only, leaving for nurture, at best, a secondary role. A larger sample of barred galaxies is needed to shed more light on the relationship between environment and the presence of bars.

According to our results, the relation between environment and bars appears to be, at best, a second order effect. Since the NA10 sample contains galaxies in a wide range of environments (see histograms in Fig. 1), it is worth emphasising the fact that the presence of bars do not seem to depend on environment while most galaxy properties do.

ACKNOWLEDGEMENTS

This work has been partially supported with grants from Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina (CONICET), Secretaría de Ciencia y Tecnología de la Universidad de Córdoba and Ministerio de Ciencia y Tecnología de la Provincia de Córdoba, Argentina.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the NSF, the U.S. Department of Energy, NASA, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>. The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Inst. Potsdam, Univ. of Basel, Univ. of Cambridge, Case Western Reserve Univ., Univ. of Chicago, Drexel Univ., Fermilab, the Inst. for Advanced Study, the Japan Participation Group, Johns Hopkins Univ., the Joint Inst. for Nuclear Astrophysics, the Kavli Inst. for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Inst. for Astronomy (MPIA), the Max-Planck-Inst. for Astrophysics (MPA), New Mexico State Univ., Ohio State Univ., Univ. of Pittsburgh, Univ. of Portsmouth, Princeton Univ., the United States Naval Observatory, and the Univ. of Washington.

REFERENCES

- Abraham, R. G., Merrifield, M. R., Ellis, R. S., Tanvir, N. R., & Brinchmann, J. 1999, *MNRAS*, 308, 569
- Adelman-McCarthy J. K., et al. 2006, *ApJS*, 162, 38
- Athanassoula, E. 2003, *MNRAS*, 341, 1179
- Athanassoula, E. 2005, *Celestial Mechanics and Dynamical Astronomy*, 91, 9
- Athanassoula, E. 2008, *MNRAS*, 390, L69
- Baldray, I. K., Balogh, M. L., Bower, R. G., Glazebrook, K., Nichol, R. C., Bamford, S. P., & Budavari, T. 2006, *MNRAS*, 373, 469
- Berentzen, I., Athanassoula, E., Heller, C. H., & Fricke, K. J. 2004, *MNRAS*, 347, 220
- Blanton, M. R., et al. 2003, *ApJ*, 594, 186
- Blanton, M. R., Eisenstein, D., Hogg, D. W., Schlegel, D. J., & Brinkmann, J. 2005, *ApJ*, 629, 143
- Brinchmann, J., Charlot, S., White, S. D. M., Tremonti, C., Kauffmann, G., Heckman, T., & Brinkmann, J. 2004, *MNRAS*, 351, 1151
- Buta, R. 1986, *ApJS*, 61, 609
- Buta, R., & Combes, F. 1996, *FCP*, 17, 95
- Corsini, E. M., Debattista, V. P., & Aguerri, J. A. L. 2003, *ApJL*, 599, L29
- Courteau, S., Andersen, D. R., Bershad, M. A., MacArthur, L. A., & Rix, H.-W. 2003, *ApJ*, 594, 208
- de Vaucouleurs, G. 1963, *ApJS*, 8, 31
- Debattista, V. P., & Sellwood, J. A. 1998, *ApJL*, 493, L5
- Elmegreen, B. G., & Elmegreen, D. M. 1985, *ApJ*, 288, 438
- Elmegreen, D. M., Elmegreen, B. G., & Bellin, A. D. 1990, *ApJ*, 364, 415
- Elmegreen, B. G., Elmegreen, D. M., Knapen, J. H., Buta, R. J., Block, D. L., & Puerari, I. 2007, *ApJL*, 670, L97
- Erwin, P. 2005, *MNRAS*, 364, 283
- Eskridge, P. B., et al. 2000, *AJ*, 119, 536
- Gerin, M., Combes, F., & Athanassoula, E. 1990, *A&A*, 230, 37
- Heller, C. H., Shlosman, I., & Athanassoula, E. 2007, *ApJ*, 671, 226
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997, *ApJ*, 487, 591
- Kauffmann, G., et al. 2003b, *MNRAS*, 341, 54
- Kenney, J. D. P., & Lord, S. D. 1991, *ApJ*, 381, 118
- Kormendy, J. 1982, *ApJ*, 257, 75
- Kormendy, J., & Kennicutt, R. C., Jr. 2004, *ARAA*, 42, 603
- Laurikainen, E., Salo, H., & Buta, R. 2004, *ApJ*, 607, 103
- Lindblad, P. O. 1960, *Stockholms Observatoriums Annaler*, 21, 4
- Martin, P., & Roy, J.-R. 1994, *ApJ*, 424, 599
- Martínez, H. J., & Muriel, H. 2006, *MNRAS*, 370, 1003
- Martínez, H. J., Coenda, V., & Muriel, H. 2008, *MNRAS*, 391, 585
- Méndez-Abreu, J., Sánchez-Janssen, R., & Aguerri, J. A. L. 2010, *ApJL*, 711, L61
- Menéndez-Delmestre, K., Sheth, K., Schinnerer, E., Jarrett, T. H., & Scoville, N. Z. 2007, *ApJ*, 657, 790
- Mihos, J. C., McGaugh, S. S., & de Blok, W. J. G. 1997, *ApJL*, 477, L79
- Miwa, T., & Noguchi, M. 1998, *ApJ*, 499, 149
- Mulchaey, J. S., & Regan, M. W. 1997, *ApJL*, 482, L135
- Nair, P. B., & Abraham, R. G. 2010a, *ApJS*, 186, 427
- Nair, P. B., & Abraham, R. G. 2010b, *ApJL*, 714, L260
- Noguchi, M. 1987, *MNRAS*, 228, 635
- Schmidt, M. 1968, *ApJ*, 151, 393
- Schwarz, M. P. 1981, *ApJ*, 247, 77
- Sellwood, J. A., & Wilkinson, A. 1993, *Reports on Progress in Physics*, 56, 173
- Sérsic, J. L., & Pastoriza, M. 1967, *PASP*, 79, 152
- Shlosman, I., Begelman, M. C., & Frank, J. 1990, *Nature*, 345, 679
- Sheth, K., Regan, M. W., Vogel, S. N., & Teuben, P. J. 2000, *ApJ*, 532, 221
- Sheth, K., Vogel, S. N., Regan, M. W., Teuben, P. J., Harris, A. I., & Thornley, M. D. 2002, *AJ*, 124, 2581
- Sheth, K., Vogel, S. N., Regan, M. W., Thornley, M. D., & Teuben, P. J. 2005, *ApJ*, 632, 217
- Sheth, K., et al. 2008, *ApJ*, 675, 1141
- Strauss, M. A., et al. 2002, *AJ*, 124, 1810
- van den Bergh, S., Abraham, R. G., Ellis, R. S., Tanvir, N. R., Santiago, B. X., & Glazebrook, K. G. 1996, *AJ*, 112, 359
- van den Bergh, S. 2002, *AJ*, 124, 782
- Walker, I. R., Mihos, J. C., Hernquist, L., Bolte, M., & Mendes de Oliveira, C. 1996, *IAU Colloq. 157: Barred Galaxies*, 91, 486
- Weinberg, M. D. 1985, *MNRAS*, 213, 451
- Yang, X., Mo, H. J., van den Bosch, F. C., Pasquali, A., Li, C., & Barden, M. 2007, *ApJ*, 671, 153
- Zaritsky, D. 1992, *ApJL*, 390, L73